

COMBINATORIAL IDENTITIES INVOLVING MERTENS FUNCTION THROUGH RELATIVELY PRIME SUBSETS

MOHAMED EL BACHRAOUI

ABSTRACT. In this note we give some identities which involve the Mertens function $M(n)$. Our proofs are combinatorial with relatively prime subsets as a main tool.

1. INTRODUCTION

Mertens function given by

$$M(n) = \sum_{d=1}^n \mu(d),$$

where μ denotes the Möbius mu function, is an important function in (analytic) number theory. Most of mathematical identities where $M(n)$ appears are either recursive formulas for $M(n)$ or formulas with an analytic flavor relating $M(n)$ to other functions. For a survey on identities involving $M(n)$ we refer to [1, 4] and for a survey on combinatorial identities involving other arithmetical functions we refer to [7, 10] and their references. In this work we will give some other identities involving the function $M(n)$. Our proofs are combinatorial and based on relatively prime subsets of sets of positive integers. We list two of the identities which we intend to prove. Let $\lfloor x \rfloor$ denote the floor of x .

1) If $n > 3$, then

$$\sum_{d=1}^n \mu(d) 2^{\lfloor \frac{n}{d} \rfloor - \lfloor \frac{n-3}{d} \rfloor} = \begin{cases} 3 + M(n), & \text{if } n \text{ is even} \\ 4 + M(n), & \text{if } n \text{ is odd.} \end{cases}$$

2) If $1 < m < n$, then

$$\sum_{d=1}^{n+1} \mu(d) 2^{\lfloor \frac{n+1}{d} \rfloor - \lfloor \frac{n-1}{d} \rfloor + \lfloor \frac{m}{d} \rfloor - \lfloor \frac{m-1}{d} \rfloor} = \begin{cases} 2 + M(n+1), & \text{if } (m, n) > 1 \text{ and } (m, n+1) > 1 \\ 3 + M(n+1), & \text{if } (m, n) = 1 \text{ and } (m, n+1) > 1 \text{ or } (m, n) > 1 \text{ and } (m, n+1) = 1 \\ 4 + M(n+1), & \text{if } (m, n) = (m, n+1) = 1. \end{cases}$$

Date: December 8, 2009.

1991 Mathematics Subject Classification. 11A25, 11B05, 11B75.

Key words and phrases. Combinatorial identities, Mertens function, Möbius function, Relatively prime sets.

2. RELATIVELY PRIME SUBSETS OF $[l_1, m_1] \cup [l_2, m_2]$

Throughout this section let k, l, m, l_1, l_2, m_1 , and m_2 be positive integers such that $l \leq m$, $l_1 \leq m_1$ and $l_2 \leq m_2$, let $[l, m] = \{l, l+1, \dots, m\}$, and let A be a nonempty finite set of positive integers. The set A is called *relatively prime* if $\gcd(A) = 1$.

Definition 1. *Let*

$$f(A) = \#\{X \subseteq A : X \neq \emptyset \text{ and } \gcd(X) = 1\}$$

and

$$f_k(A) = \#\{X \subseteq A : \#X = k \text{ and } \gcd(X) = 1\}.$$

Nathanson in [8] introduced among other functions $f(n)$ and $f_k(n)$ (in our terminology $f([1, n])$ and $f_k([1, n])$ respectively) and found

$$(1) \quad f([1, n]) = \sum_{d=1}^n \mu(d)(2^{\lfloor \frac{n}{d} \rfloor} - 1) \text{ and } f_k([1, n]) = \sum_{d=1}^n \mu(d) \binom{\lfloor \frac{n}{d} \rfloor}{k}.$$

Formulas for $f([m, n])$ and $f_k([m, n])$ are found in [5, 9]. Recently Ayad and Kihel in [3] considered relatively prime subsets of sets which are in arithmetic progression and obtained formulas for $f([l, m])$ and $f_k([l, m])$ as consequences since the integer interval $[l, m]$ is in arithmetic progression. However the authors' argument seems not to extend to unions of integer intervals. In this section we will give formulas for $f([l_1, m_1] \cup [l_2, m_2])$ and for $f_k([l_1, m_1] \cup [l_2, m_2])$. For the sake of completeness we include the following result which is a natural extension of [5, Theorem 2 (a)] on Möbius inversion for arithmetical functions of several variables. For simplicity of notation we let

$$(\overline{m}_a, \overline{n}_b) = (m_1, m_2, \dots, m_a, n_1, n_2, \dots, n_b)$$

and

$$\left(\frac{\overline{m}_a}{d}, \left\lfloor \frac{\overline{n}_b}{d} \right\rfloor \right) = \left(\frac{m_1}{d}, \frac{m_2}{d}, \dots, \frac{m_a}{d}, \left\lfloor \frac{n_1}{d} \right\rfloor, \left\lfloor \frac{n_2}{d} \right\rfloor, \dots, \left\lfloor \frac{n_b}{d} \right\rfloor \right).$$

Theorem 2. *If F and G are arithmetical of $a + b$ variables, then*

$$G(\overline{m}_a, \overline{n}_b) = \sum_{d|(m_1, m_2, \dots, m_a)} F\left(\frac{\overline{m}_a}{d}, \left\lfloor \frac{\overline{n}_b}{d} \right\rfloor\right)$$

if and only if

$$F(\overline{m}_a, \overline{n}_b) = \sum_{d|(m_1, m_2, \dots, m_a)} \mu(d) G\left(\frac{\overline{m}_a}{d}, \left\lfloor \frac{\overline{n}_b}{d} \right\rfloor\right).$$

We need the following three lemmas the proofs of which can be obtained using the same sort of idea and therefore we prove only the first one.

Lemma 3. *Let*

$$g(m_1, l_2, m_2) = \#\{X \subseteq [1, m_1] \cup [l_2, m_2] : l_2 \in X \text{ and } \gcd(X) = 1\},$$

$$g_k(m_1, l_2, m_2) = \#\{X \subseteq [1, m_1] \cup [l_2, m_2] : l_2 \in X, |X| = k, \text{ and } \gcd(X) = 1\}.$$

Then

$$(a) \quad g(m_1, l_2, m_2) = \sum_{d|l_2} \mu(d) 2^{\lfloor m_1/d \rfloor + \lfloor m_2/d \rfloor - l_2/d},$$

$$(b) \quad g_k(m_1, l_2, m_2) = \sum_{d|l_2} \mu(d) \binom{\lfloor m_1/d \rfloor + \lfloor m_2/d \rfloor - l_2/d}{k-1}.$$

Proof. (a) Let $\mathcal{P}(m_1, l_2, m_2)$ denote the set of subsets of $[1, m_1] \cup [l_2, m_2]$ containing l_2 and let $\mathcal{P}(m_1, l_2, m_2, d)$ be the set of subsets X of $[1, m_1] \cup [l_2, m_2]$ such that $l_2 \in X$ and $\gcd(X) = d$. It is clear that the set $\mathcal{P}(m_1, l_2, m_2)$ of cardinality $2^{m_1+m_2-l_2}$ can be partitioned using the equivalence relation of having the same gcd (dividing l_2). Moreover, the mapping $A \mapsto \frac{1}{d}A$ is a one-to-one correspondence between $\mathcal{P}(m_1, l_2, m_2, d)$ and the set of subsets Y of $[1, \lfloor m_1/d \rfloor] \cup [l_2/d, \lfloor m_2/d \rfloor]$ such that $l_2/d \in Y$ and $\gcd(Y) = 1$. Then

$$2^{m_1+m_2-l_2} = \sum_{d|l_2} \#\mathcal{P}(m_1, l_2, m_2, d) = \sum_{d|l_2} g(\lfloor m_1/d \rfloor, l_2/d, \lfloor m_2/d \rfloor),$$

which by Theorem 2 is equivalent to

$$g(m_1, l_2, m_2) = \sum_{d|l_2} \mu(d) 2^{\lfloor m_1/d \rfloor + \lfloor m_2/d \rfloor - l_2/d}.$$

(b) Similarly

$$\binom{m_1 + m_2 - l_2}{k-1} = \sum_{d|l_2} g_k(\lfloor m_1/d \rfloor, l_2/d, \lfloor m_2/d \rfloor),$$

which by Theorem 2 is equivalent to

$$g_k(m_1, l_2, m_2) = \sum_{d|l_2} \mu(d) \binom{\lfloor m_1/d \rfloor + \lfloor m_2/d \rfloor - l_2/d}{k-1}.$$

□

Lemma 4. *Let*

$$h^{(1)}(l_1, m_1) = \#\{X \subseteq [l_1, m_1] : l_1 \in X \text{ and } \gcd(X) = 1\},$$

$$h_k^{(1)}(l_1, m_1) = \#\{X \subseteq [l_1, m_1] : l_1 \in X, \#X = k, \text{ and } \gcd(X) = 1\}.$$

Then

$$(a) \quad h^{(1)}(l_1, m_1) = \sum_{d|l_1} \mu(d) 2^{\lfloor m_1/d \rfloor - l_1/d},$$

$$(b) \quad h_k^{(1)}(l_1, m_1) = \sum_{d|l_1} \mu(d) \binom{\lfloor m_1/d \rfloor - l_1/d}{k}.$$

Lemma 5. *Let*

$$h^{(2)}(l_1, m_1, l_2, m_2) = \#\{X \subseteq [l_1, m_1] \cup [l_2, m_2] : l_1, l_2 \in X \text{ and } \gcd(X) = 1\},$$

$$h_k^{(2)}(l_1, m_1, l_2, m_2) = \#\{X \subseteq [l_1, m_1] \cup [l_2, m_2] : l_1, l_2 \in X, |X| = k, \text{ and } \gcd(X) = 1\}.$$

Then

$$(a) \quad h^{(2)}(l_1, m_1, l_2, m_2) = \sum_{d|(l_1, l_2)} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{l_1+l_2}{d}},$$

$$(b) \quad h_k^{(2)}(m_1, l_2, m_2) = \sum_{d|(l_1, l_2)} \mu(d) \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{l_1+l_2}{d}}{k-2}.$$

We further need the following special case to prove the main theorem of this section.

Theorem 6. *We have*

$$(a) \quad f([1, m_1] \cup [l_2, m_2]) = \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor} - 1),$$

$$(b) \quad f_k([1, m_1] \cup [l_2, m_2]) = \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor}{k}.$$

Proof. (a) Clearly

$$\begin{aligned} f([1, m_1] \cup [l_2, m_2]) &= f([1, m_2]) - \sum_{i=m_1+1}^{l_2-1} g(m_1, i, m_2) \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor m_2/d \rfloor} - 1) - \sum_{i=m_1+1}^{l_2-1} \sum_{d|i} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{i}{d}} \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor m_2/d \rfloor} - 1) - \sum_{d=1}^{l_2-1} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor} \sum_{j=\lfloor \frac{m_1}{d} \rfloor + 1}^{\lfloor \frac{l_2-1}{d} \rfloor} 2^{-j} \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor m_2/d \rfloor} - 1) - \sum_{d=1}^{l_2-1} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor} 2^{-\lfloor \frac{m_1}{d} \rfloor} \left(1 - 2^{-\lfloor \frac{l_2-1}{d} \rfloor + \lfloor \frac{m_1}{d} \rfloor}\right) \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor m_2/d \rfloor} - 1) - \sum_{d=1}^{m_2} \mu(d) 2^{\lfloor \frac{m_2}{d} \rfloor} \left(1 - 2^{-\lfloor \frac{l_2-1}{d} \rfloor + \lfloor \frac{m_1}{d} \rfloor}\right) \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor} - 1), \end{aligned}$$

where the second identity follows by (1) and Lemma 3.

(b) We have

$$\begin{aligned} f_k([1, m_1] \cup [l_2, m_2]) &= f_k([1, m_2]) - \sum_{i=m_1+1}^{l_2-1} g(m_1, i, m_2) \\ &= \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor m_2/d \rfloor}{k} - \sum_{i=m_1+1}^{l_2-1} \sum_{d|i} \mu(d) \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{i}{d}}{k-1} \\ &= \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor m_2/d \rfloor}{k} - \sum_{d=1}^{m_2} \mu(d) \sum_{j=\lfloor \frac{m_1}{d} \rfloor + 1}^{\lfloor \frac{l_2-1}{d} \rfloor} \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{i}{d}}{k-1} \\ &= \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor m_2/d \rfloor}{k} - \sum_{d=1}^{m_2} \mu(d) \sum_{i=\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{l_2-1}{d}}^{\lfloor \frac{m_2}{d} \rfloor - 1} \binom{i}{k-1} \\ &= \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor m_2/d \rfloor}{k} - \sum_{d=1}^{m_2} \mu(d) \left(\binom{\lfloor m_2/d \rfloor}{k} - \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{l_2-1}{d}}{k} \right) \\ &= \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{l_2-1}{d}}{k}. \end{aligned}$$

This completes the proof. $\square$

We are now ready to prove the main theorem of this section.

Theorem 7. *We have*

$$(a) \quad f([l_1, m_1] \cup [l_2, m_2]) = \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_1-1}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor} - 1),$$

$$(b) \quad f_k([l_1, m_1] \cup [l_2, m_2]) = \sum_{d=1}^{m_2} \mu(d) \binom{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_1-1}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor}{k}.$$

Proof. (a) Clearly

$$(3) \quad \begin{aligned} f([l_1, m_1] \cup [l_2, m_2]) &= f([1, m_1] \cup [l_2, m_2]) - \sum_{i=1}^{l_1-1} \sum_{j=l_2}^{m_2} h^{(2)}(i, m_1, j, m_2) - \sum_{i=1}^{l_1-1} h^{(1)}(i, m_1) \\ &= \sum_{d=1}^{m_2} \mu(d) (2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor} - 1) - \sum_{i=1}^{l_1-1} \sum_{j=l_2}^{m_2} \sum_{d|(i,j)} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{i+j}{d}} - \sum_{i=1}^{l_1-1} \sum_{d|i} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor - \frac{i}{d}}, \end{aligned}$$

where the second identity follows by Theorem 6, Lemma 4, and Lemma 5. Rearranging the triple summation in identity (3), we get

$$(4) \quad \begin{aligned} \sum_{i=1}^{l_1-1} \sum_{j=l_2}^{m_2} \sum_{d|(i,j)} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \frac{i+j}{d}} &= \sum_{d=1}^{m_2} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor} \sum_{i=1}^{\lfloor \frac{l_1-1}{d} \rfloor} 2^{-i} \sum_{j=\lfloor \frac{l_2-1}{d} \rfloor + 1}^{\lfloor \frac{m_2}{d} \rfloor} 2^{-j} \\ &= \sum_{d=1}^{m_2} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor} (1 - 2^{-\lfloor \frac{m_2}{d} \rfloor + \lfloor \frac{l_2-1}{d} \rfloor}) (1 - 2^{-\lfloor \frac{l_1-1}{d} \rfloor}). \end{aligned}$$

Similarly the last double summation in identity (3) gives

$$(5) \quad \begin{aligned} \sum_{i=1}^{l_1-1} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor} \sum_{i=1}^{\lfloor \frac{l_1-1}{d} \rfloor} 2^{-\frac{i}{d}} &= \sum_{d=1}^{l_1-1} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor} (1 - 2^{-\lfloor \frac{l_1-1}{d} \rfloor}) \\ &= \sum_{d=1}^{m_2} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor} (1 - 2^{-\lfloor \frac{l_1-1}{d} \rfloor}). \end{aligned}$$

Appealing to identities (3), (4), and (5) we find

$$f([l_1, m_1] \cup [l_2, m_2]) = \sum_{d|n} \mu(d) 2^{\lfloor \frac{m_1}{d} \rfloor + \lfloor \frac{m_2}{d} \rfloor - \lfloor \frac{l_1-1}{d} \rfloor - \lfloor \frac{l_2-1}{d} \rfloor}.$$

This completes the proof of part (a). Part (b) follows by similar arguments. $\square$

Corollary 8. (Ayad-Kihel [3]) *We have*

$$(a) \quad f([l, m]) = \sum_{d=1}^m (2^{\lfloor \frac{m}{d} \rfloor - \lfloor \frac{l-1}{d} \rfloor} - 1),$$

$$(b) \quad f_k([l, m]) = \sum_{d=1}^m \binom{\lfloor \frac{m}{d} \rfloor - \lfloor \frac{l-1}{d} \rfloor}{k}.$$

Proof. Use Theorem 7 with $l_1 = l$, $m_1 = m - 1$, and $l_2 = m_2 = m$. $\square$

3. COMBINATORIAL IDENTITIES

Theorem 9. *If $n > 1$, then*

$$\sum_{d=1}^{n+1} \mu(d) 2^{\lfloor \frac{n+1}{d} \rfloor - \lfloor \frac{n-1}{d} \rfloor} = 1 + M(n+1)$$

Proof. Apply Corollary 8 to the interval $[n, n+1]$ and use the obvious fact that $f([n, n+1]) = 1$. $\square$

Theorem 10. *In $n > 3$, then*

$$\sum_{d=1}^n 2^{\lfloor \frac{n}{d} \rfloor - \lfloor \frac{n-3}{d} \rfloor} = \begin{cases} 3 + M(n), & \text{if } n \text{ is even} \\ 4 + M(n), & \text{if } n \text{ is odd.} \end{cases}$$

Proof. Combine Corollary 8 applied to the interval $[n-2, n]$ and the fact that

$$f([n-2, n]) = \begin{cases} 3, & \text{if } n \text{ is even} \\ 4, & \text{if } n \text{ is odd.} \end{cases}$$

$\square$

Theorem 11. (a) *If $1 < m < n$, then*

$$\begin{aligned} & \sum_{d=1}^{n+1} \mu(d) 2^{\lfloor \frac{n+1}{d} \rfloor - \lfloor \frac{n-1}{d} \rfloor + \lfloor \frac{m}{d} \rfloor - \lfloor \frac{m-1}{d} \rfloor} = \\ & \begin{cases} 2 + M(n+1), & \text{if } (m, n) > 1 \text{ and } (m, n+1) > 1 \\ 3 + M(n+1), & \text{if } (m, n) = 1 \text{ and } (m, n+1) > 1 \text{ or } (m, n) > 1 \text{ and } (m, n+1) = 1 \\ 4 + M(n+1), & \text{if } (m, n) = (m, n+1) = 1. \end{cases} \end{aligned}$$

(b) *If $1 < n < m-1$, then*

$$\begin{aligned} & \sum_{d=1}^m \mu(d) 2^{\lfloor \frac{n+1}{d} \rfloor - \lfloor \frac{n-1}{d} \rfloor + \lfloor \frac{m}{d} \rfloor - \lfloor \frac{m-1}{d} \rfloor} = \\ & \begin{cases} 2 + M(m), & \text{if } (m, n) > 1 \text{ and } (m, n+1) > 1 \\ 3 + M(m), & \text{if } (m, n) = 1 \text{ and } (m, n+1) > 1 \text{ or } (m, n) > 1 \text{ and } (m, n+1) = 1 \\ 4 + M(m), & \text{if } (m, n) = (m, n+1) = 1. \end{cases} \end{aligned}$$

Proof. (a) Clearly

$$f([m, m] \cup [n, n+1]) = \begin{cases} 2, & \text{if } (m, n) > 1 \text{ and } (m, n+1) > 1 \\ 3, & \text{if } (m, n) = 1 \text{ and } (m, n+1) > 1 \text{ or } (m, n) > 1 \text{ and } (m, n+1) = 1 \\ 4, & \text{if } (m, n) = (m, n+1) = 1. \end{cases}$$

Combine this identity with Theorem 7(a) applied to $[m, m] \cup [n, n+1]$.

(b) Similar to part (a) with an application of Theorem 7(a) to $[n, n+1] \cup [m, m]$. $\square$

REFERENCES

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, Undergraduate Texts in Mathematics, Springer, 1 edition, 1976.
- [2] Mohamed Ayad and Omar Kihel, *On the Number of Subsets Relatively Prime to an Integer*, Journal of Integer Sequences, **Vol. 11**, (2008), Article 08.5.5.
- [3] Mohamed Ayad and Omar Kihel, *On Relatively Prime Sets*, Integers **9**, (2009), 343-352.
- [4] F. Dress, *Fonction sommatoire de la fonction de Möbius. I. Majorations expérimentales*, Experiment. Math. **2**, (1993), 89-98.
- [5] Mohamed El Bachraoui, *The number of relatively prime subsets and phi functions for sets $\{m, m+1, \dots, n\}$* , Integers **7** (2007), A43, 8pp.
- [6] Mohamed El Bachraoui, *On the Number of Subsets of $[1, m]$ Relatively Prime to n and Asymptotic Estimates*, Integers **8** (2008), A 41, 5 pp.
- [7] M. Hall, *Combinatorial Theory*, Second edition, Wiley, New York, 1986.
- [8] Melvyn B. Nathanson, *Affine invariants, relatively prime sets, and a phi function for subsets of $\{1, 2, \dots, n\}$* , Integers **7** (2007), A1, 7pp.
- [9] Melvyn B. Nathanson and Brooke Orosz, *Asymptotic estimates for phi functions for subsets of $\{m+1, m+2, \dots, n\}$* , Integers **7** (2007), A54, 5pp.
- [10] J. Riordan, *Combinatorial Identities*, Wiley, New York, 1968.

DEPT. MATH. SCI, UNITED ARAB EMIRATES UNIVERSITY, PO BOX 17551, AL-AIN, UAE
E-mail address: melbachraoui@uaeu.ac.ae